

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/313918596>

Medical muddles that maim our children with allergies, asthma and autism

Technical Report · February 2017

DOI: 10.13140/RG.2.2.13347.45603

CITATIONS

0

READS

1,354

1 author:



Vinu Arumugham

21 PUBLICATIONS 18 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Vaccine Safety Analysis [View project](#)

All content following this page was uploaded by [Vinu Arumugham](#) on 24 February 2017.

The user has requested enhancement of the downloaded file.

Medical muddles that maim our children with allergies, asthma and autism

Introduction

In the 1900s it was discovered that injected allergens cause the development of allergy to that allergen.¹⁻³

In 1911, Wells and Osborne⁴ discovered that eating allergens, protected against developing allergy upon subsequent allergen injection.

Researchers confuse allergen injection with allergen ingestion

Grimshaw et al.⁵ recently described a 1906 concept of allergen priming. That is, the development of an allergy upon first exposure to the allergen. They describe it as the basis for researchers concluding that the first exposure of allergen in utero and breastfeeding causes the development of food allergies. This led to the guidance (in the 80s) of avoiding dietary allergen exposure in pregnant and lactating women. The guidance also delayed dietary allergen exposure in infants.⁵

MISTAKE 1: The 1906 concept of allergen priming was caused by first exposure to the allergen via **injection**, not by in utero, breastfeeding or dietary exposure.

This mistake had two consequences.

- a) The real cause of allergy - injection of allergens (via food protein contaminated vaccines), continued unchanged.
- b) The protection children got from eating allergens early in life, was lost due to the guidance delaying allergen exposure.

The combined effect of this mistake was a **further increase in food allergy prevalence**.

The guidance of allergen avoidance during pregnancy/lactation has now been withdrawn.⁵ The guidance to introduce allergens early in infants is being rolled out now.⁶ But these steps only fix one half of the problem created above.

MISTAKE 2: Allergens are still continuing to be injected into children, ignoring the 1906 finding.

The result is an epidemic of life-threatening food allergies, asthma and life-ruining autism, as detailed in the articles below.

Food Allergy

[Professional Misconduct by NAM Committee on Food Allergy](#)

Autism

[Autism Spectrum Disorders: A special case of vaccine-induced cow's milk allergy?](#)

Vaccines causing food allergies and asthma is established science

Food protein contaminated vaccines causing the development of food allergies⁷ and asthma is **established science**. Every medical student is taught this basic fact.

Medical Immunology notes from the University of California, Irvine, School of Medicine.
<http://jeeves.mmg.uci.edu/immunology/CoreNotes/Chap21.pdf>

pg. 157:

“A guinea pig can be sensitized by intramuscular injection of an antigen, say OVA (ovalbumin). Its immune system responds by producing antibody to OVA, including (but not exclusively) IgE. Some of this circulating IgE will be fixed onto mast cells in various tissues, including the vasculature and respiratory tract. Three weeks later, the same animal can be challenged either with an intravenous dose of OVA or by exposure to an aerosol containing OVA. Following IV injection, the animal will rapidly develop severe vascular shock and die within a few minutes (the combination of venule constriction and capillary dilation results in pooling of blood in the peripheral circulation and a drastic drop in blood pressure). If exposed to the aerosol, it will equally rapidly die from bronchial constriction, an experimental model for human asthma.”

The guinea pig is considered the "experimental model for human asthma" because it is **representative** of human body behavior.

Doctors today inject numerous food protein contaminated vaccines into babies. When they predictably develop food allergies and asthma, the doctors **inexplicably** claim, they **don't know** what caused these diseases! And inexplicably, **do not** report this to the Vaccine Adverse Event Reporting System (VAERS).

Allergens in vaccines per the National Academy of Medicine (NAM) report

<https://www.nap.edu/catalog/23658/finding-a-path-to-safety-in-food-allergy-assessment-of>

NAM report pg.241

"Allergens in Vaccines, Medications, and Dietary Supplements

Physicians and patients with food allergy must consider potential food allergen exposures in vaccines, medications, and dietary supplement products (e.g., vitamins, probiotics), which are not regulated by labelling laws. Also, excipients (i.e., substances added to medications to improve various characteristics) may be food or derived from foods (Kelso, 2014). These

include milk proteins; soy derivatives; oils from sesame, peanut, fish or soy; and beef or fish gelatin. The medications involved include vaccines; anesthetics; and oral, topical, and injected medications. With perhaps the exception of gelatin, reactions appear to be rare overall, likely because little residual protein is included in the final preparation of these items. The specific risk for each medication is not known.

Vaccines also may contain food allergens, such as egg protein or gelatin."

Vaccines are a cause of asthma (may not be the only cause)

There are at least two mechanisms by which vaccines can cause asthma.

As in the guinea pig model above, people get sensitized (IgE mediated) by food protein injections. Upon inhalation of aerosolized food particles, they can suffer asthma symptoms.

Again, as in the guinea pig model, people can get sensitized to pathogen associated vaccine antigens (IgE mediated).⁸⁻¹³ Upon inhalation of aerosolized pathogen particles, (influenza viral particles, pertussis bacterial particles, meningococcal bacterial particles, yeast, etc.) they will develop asthma symptoms.

The IOM 2011¹⁴ report on vaccine adverse events **wrongly** concluded that influenza vaccines do not cause asthma. There are a couple of reasons.

1. For the above mentioned asthma mechanism, one has to wait a few weeks after the vaccine for IgE mediated sensitization to occur.
2. For this asthma mechanism, patients must be exposed to aerosolized food proteins or aerosolized pathogen proteins to trigger asthma symptoms.

The studies the IOM reviewed **did not** address the above two conditions. Thus, the IOM's conclusion on this matter is wrong.

Those who don't learn from history ...

Silverstein¹ describes medical discoveries being forgotten and rediscovered. Not only that, we are finding out that discoveries are misunderstood, misinterpreted and ignored, with devastating consequences.

Action

All contaminating proteins in vaccines must be removed immediately.¹⁵ All proteins in vaccines needed for disease protection, must also be thoroughly scrutinized for molecular mimicry that can result in allergic or autoimmune diseases.

References

1. Silverstein AM. Clemens Freiherr von Pirquet: Explaining immune complex disease in 1906. *Nat Immunol.* 2000 Dec;1(6):453–5.
2. Wells HG. Studies on the Chemistry of Anaphylaxis. *J Infect Dis.* Oxford University Press; 1908;5(4):449–83.
3. Wells HG. Studies on the Chemistry of Anaphylaxis (III). Experiments with Isolated Proteins, Especially Those of the Hen's Egg. *J Infect Dis.* Oxford University Press; 1911;9(2):147–71.
4. Wells HG, Osborne TB. The Biological Reactions of the Vegetable Proteins I. Anaphylaxis. *J Infect Dis.* 1911;8(1):66–124.
5. Grimshaw K, Logan K, O'Connell S, Kiely M, Patient K, van Bilsen J, et al. Modifying the infant's diet to prevent food allergy. *Arch Dis Child.* 2017 Jan 18;102(2):179 LP – 186.
6. Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med.* 2015;372(9):803–13.
7. Arumugham V. Evidence that Food Proteins in Vaccines Cause the Development of Food Allergies and Its Implications for Vaccine Policy. *J Dev Drugs.* 2015;4(137):2.
8. Nakayama T, Kumagai T, Nishimura N, Ozaki T, Okafuji T, Suzuki E, et al. Seasonal split influenza vaccine induced IgE sensitization against influenza vaccine. *Vaccine.* 2015 Nov 9;33(45):6099–105.
9. Davidsson A, Eriksson JC, Rudblad S, Brokstad KA. Influenza specific serum IgE is present in non-allergic subjects. *Scand J Immunol.* 2005 Dec;62(6):560–1.
10. Ryan EJ, Nilsson L, Kjellman NIM, Gothefors L, Mills KHG. Booster immunization of children with an acellular pertussis vaccine enhances Th2 cytokine production and serum IgE responses against pertussis toxin but not against common allergens. *Clin Exp Immunol.* 2000;121(2):193–200.
11. Holt PG, Snelling T, White OJ, Sly PD, DeKlerk N, Carapetis J, et al. Transiently increased IgE responses in infants and pre-schoolers receiving only acellular Diphtheria–Pertussis–Tetanus (DTaP) vaccines compared to those initially receiving at least one dose of cellular vaccine (DTwP) – Immunological curiosity or canary i. *Vaccine.* 2016 Jul 29;34(35):4257–62.
12. Smith-Norowitz T a, Wong D, Kusunruksa M, Norowitz KB, Joks R, Durkin HG, et al. Long term persistence of IgE anti-influenza virus antibodies in pediatric and adult serum post vaccination with influenza virus vaccine. *Int J Med Sci.* 2011;8(3):239–44.
13. Nenoff P, Müller B, Sander U, Kunze G, Bröker M, Haustein U-F. IgG and IgE immune response against the surface glycoprotein gp200 of *Saccharomyces cerevisiae* in patients with atopic dermatitis. *Mycopathologia.* 2001;152(1):15–21.

14. Clayton EW, Rusch E, Ford A, Stratton K. *Adverse Effects of Vaccines: Evidence and Causality*. National Academies Press; 2012.
15. Kuno-Sakai H, Kimura M. Removal of gelatin from live vaccines and DTaP—an ultimate solution for vaccine-related gelatin allergy. *Biologicals*. 2003;31(4):245–9.